



Collision-dependent power law scalings in two dimensional gyrokinetic turbulence

S. S. Cerri, A. Bañón Navarro, F. Jenko, and D. Told

Citation: *Physics of Plasmas* (1994-present) **21**, 082305 (2014); doi: 10.1063/1.4892347

View online: <http://dx.doi.org/10.1063/1.4892347>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/pop/21/8?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[Alfvén wave collisions, the fundamental building block of plasma turbulence. IV. Laboratory experiment](#)
Phys. Plasmas **20**, 072901 (2013); 10.1063/1.4813242

[Alfvén wave collisions, the fundamental building block of plasma turbulence. II. Numerical solution](#)
Phys. Plasmas **20**, 072303 (2013); 10.1063/1.4812807

[Experimental study of parametric dependence of electron-scale turbulence in a spherical tokamaka\)](#)
Phys. Plasmas **19**, 056125 (2012); 10.1063/1.4719689

[Random walk of magnetic field lines in dynamical turbulence: A field line tracing method. II. Two-dimensional turbulence](#)
Phys. Plasmas **19**, 032902 (2012); 10.1063/1.3693384

[Kinetic cascade beyond magnetohydrodynamics of solar wind turbulence in two-dimensional hybrid simulations](#)
Phys. Plasmas **19**, 022305 (2012); 10.1063/1.3682960



AIP | Journal of
Applied Physics

Journal of Applied Physics is pleased to
announce **André Anders** as its new Editor-in-Chief

Collision-dependent power law scalings in two dimensional gyrokinetic turbulence

S. S. Cerri,^{1,a)} A. Bañón Navarro,¹ F. Jenko,^{1,2} and D. Told¹

¹Max-Planck-Institut für Plasmaphysik, Boltzmannstr. 2, D-85748 Garching, Germany

²Max-Planck/Princeton Center for Plasma Physics

(Received 16 May 2014; accepted 24 July 2014; published online 6 August 2014)

Nonlinear gyrokinetics provides a suitable framework to describe short-wavelength turbulence in magnetized laboratory and astrophysical plasmas. In the electrostatic limit, this system is known to exhibit a free energy cascade towards small scales in (perpendicular) real and/or velocity space. The dissipation of free energy is always due to collisions (no matter how weak the collisionality), but may be spread out across a wide range of scales. Here, we focus on freely decaying two dimensional electrostatic turbulence on sub-ion-gyroradius scales. An existing scaling theory for the turbulent cascade in the weakly collisional limit is generalized to the moderately collisional regime. In this context, non-universal power law scalings due to multiscale dissipation are predicted, and this prediction is confirmed by means of direct numerical simulations. © 2014 AIP Publishing LLC. [<http://dx.doi.org/10.1063/1.4892347>]

I. INTRODUCTION

Plasma turbulence in weakly collisional magnetized plasmas plays an important role in various systems, such as fusion devices and many space and astrophysical situations, where it leads, e.g., to anomalous transport effects and particle heating. Such plasmas are usually almost collisionless, and thus, the turbulence problem requires a kinetic approach, especially at small scales where dissipation takes place. In the standard picture, turbulence can be interpreted as a conservative transfer of energy in wavenumber space, from injection to dissipation scales.^{1,2} While the fundamental processes in the hydrodynamic case, represented by the Navier-Stokes equation, are fairly well understood at this point, the turbulence theory of magnetized plasmas is still far from complete. Over the last several years, it has become clear that plasma microturbulence—as described by nonlinear gyrokinetic (GK) theory^{3–5}—cannot be viewed as a straightforward extension of fluid turbulence.

In the 3D Navier-Stokes system, the *kinetic energy*, assumed to be injected into the system at large scales through mechanical forces, is conserved by the advective nonlinearity, which is responsible for transferring the energy from the injection scales to the smallest ones (turbulent *cascade*) at which the energy is then dissipated by viscous effects.^{1,2} In the GK formalism, on the other hand, the ideal quadratic invariant is given by the *free energy*, which is subject to a phase-space cascade towards small scales in (perpendicular) real and/or velocity space.^{6,7} A Kolmogorov-like phenomenological scaling theory of GK turbulence has been developed for the weakly collisional limit.^{8,9} The respective predictions have also been confirmed via direct numerical simulations.^{10,11} The main goal of the present work is to extend this work to the moderately collisional regime, focusing, for simplicity, on freely decaying 2D electrostatic

turbulence on sub-ion-gyroradius scales. As it will turn out, our generalization predicts non-universal power law scalings due to multiscale dissipation, and this prediction is confirmed by means of direct numerical simulations.

The remainder of this paper is organized as follows. An introduction to the GK system of equations in the 2D electrostatic limit and to its global and local energy balance equations is given in Sec. II. Then, in Sec. III, we provide a brief review of the theory of nonlinear phase mixing at sub-ion-gyroradius scales as proposed in Refs. 8 and 9 (Sec. III A), and we propose a novel natural extension of such theory to the multiscale dissipation case (Sec. III B). In Sec. IV, we present the results from direct numerical simulations with the GK plasma turbulence code GENE.^{12–14} We demonstrate the validity of the assumptions made in the theory we developed in Sec. III B for increasing collisionality by comparisons with the numerical results. Moreover, in the low collisionality limit, we recover the hypothesis and the results predicted by the standard theory of Refs. 8 and 9, showing that the transition from one case to the other is continuous, as one might have expected.

II. THE SYSTEM UNDER STUDY

In the present study, the full GK system of equations is reduced to the simple scenario of a single ion species, electrostatic perturbations, slab geometry with $k_{\parallel} = 0$ (thus avoiding parallel effects such as the Landau damping and the linear phase mixing^{15,16} to be effective), and no-response electrons (which is consistent with $k_{\perp} = 0$). No gradients in the background quantities (i.e., the density n_0 , the temperature T_0 , and the magnetic field B_0) are considered. This allows us to focus on both nonlinear and collisional effects, which are the features we are primarily interested in. We are considering the total ion distribution function F to be split into a Maxwellian part, $F_0 = (n_0/\pi^{3/2}v_T^3) \exp(-v^2/v_T^2)$, $v_T = \sqrt{2T_0/m}$ being the thermal velocity, and a perturbed part,

^{a)}Electronic mail: silvio.sergio.cerri@ipp.mpg.de

F_1 , i.e., $F = F_0 + F_1$. Then, the gyrokinetic Vlasov equation for the perturbed distribution function F_1 and the gyrokinetic Poisson equation for the self-consistent electrostatic potential ϕ_1 read

$$\frac{\partial F_1}{\partial t} + \mathbf{v}_{\bar{\phi}} \cdot \nabla F_1 = \langle C^L[F_1] \rangle, \quad (1)$$

and

$$\frac{en_0}{T_0} (1 - \Gamma_0) \phi_1 = \frac{2\pi B_0}{m} \int J_0(k_{\perp} \rho) F_1 dv_{\parallel} d\mu, \quad (2)$$

where $\mathbf{v}_{\bar{\phi}} = (c/B_0)(\mathbf{e}_z \times \nabla \bar{\phi}_1)$ is the $\mathbf{E} \times \mathbf{B}$ drift due to the gyroaveraged self-consistent electrostatic potential $\bar{\phi}_1$, the background magnetic field being aligned with the z -axis, $\langle C^L[F_1] \rangle$ is a linearized collision operator, $\Gamma_0 \equiv \frac{2\pi B_0}{mn_0} \int J_0^2(k_{\perp} \rho) F_0 dv_{\parallel} d\mu$, J_0 is the Bessel function, $\rho = v_{\perp}/\Omega_c$ is the Larmor radius, and $\mu = mv_{\perp}^2/2B_0$ is the magnetic moment. Note that, in the periodic case we are using, the gyroaverage can be written as a multiplication by the Bessel function J_0 , e.g., $\bar{\phi}_1 = J_0 \phi_1$. The term $\mathbf{v}_{\bar{\phi}} \cdot \nabla F_1$ is the nonlinear term and, because of the gyroaverage of the electrostatic potential, it is responsible for the *nonlinear phase mixing* process.^{8,9}

In the absence of collisions, due to the conservative property of the nonlinear term, Eqs. (1)–(2) admit two positive definite conserved integrals.^{8,9,17} One of them is quadratic in F_1 , which is then proportional to (minus) the perturbed part of the entropy (of the gyrocenters), while the other one is proportional to the product of $\bar{\phi}_1$ and F_1 , which is usually referred to as the electrostatic energy (or polarization term)

$$\mathcal{E}_f = \int \frac{T_0 F_1^2}{2F_0} d\Lambda, \quad (3)$$

and

$$\mathcal{E}_{\phi} = \int \frac{e \bar{\phi}_1 F_1}{2} d\Lambda, \quad (4)$$

where $d\Lambda \equiv dx dy d\Theta = \pi n_0 B_0 dx dy dv_{\parallel} d\mu$ is the phase-space element. Note that, using the Poisson equation and the local approximation, the electrostatic energy can also be written as

$$\mathcal{E}_{\phi} = \int dx dy (1 - \Gamma_0) \phi_1^2. \quad (5)$$

The entropy and the electrostatic energy, together, define the free energy

$$\mathcal{E} = \mathcal{E}_f + \mathcal{E}_{\phi}. \quad (6)$$

However, when collisions are taken into account, the global energy balance equations read

$$\frac{\partial \mathcal{E}_{\{f,\phi\}}}{\partial t} = -\mathcal{C}_{\{f,\phi\}} - \mathcal{H}_{\{f,\phi\}}, \quad (7)$$

where the collisional dissipation terms are

$$\begin{aligned} \mathcal{C}_f &= - \int \frac{T_0}{F_0} F_1 \langle C^L[F_1] \rangle d\Lambda, \\ \mathcal{C}_{\phi} &= - \int e \bar{\phi}_1 \langle C^L[F_1] \rangle d\Lambda, \end{aligned}$$

while \mathcal{H}_f and \mathcal{H}_{ϕ} are extra dissipation terms due to k_{\perp} -hyperdiffusion, which will be defined later and will be negligible for the k_{\perp} -range of interest (see Sec. IV).

Going to the local energy balance equations, in the Fourier representation they are

$$\frac{\partial E_f(k)}{\partial t} = T_f(k) - C_f(k) - D_{\perp f}(k), \quad (8a)$$

$$\frac{\partial E_{\phi}(k)}{\partial t} = T_{\phi}(k) - C_{\phi}(k) - D_{\perp, \phi}(k), \quad (8b)$$

where the spectral density of entropy $E_f(k)$ is defined by

$$\begin{aligned} \mathcal{E}_f &= \int dx dy \int \frac{T_0 F_1^2}{2F_0} d\Theta \\ &= \sum_k \int \frac{T_0 |f_k|^2}{2F_0} d\Theta = \sum_k E_f(k), \end{aligned}$$

where the sum is over all the k_x and k_y . Equivalently, we define the spectral density of the entropy collisional and perpendicular dissipations, i.e., $\mathcal{C}_f \equiv \sum_k C_f(k)$ and $\mathcal{H}_f \equiv \sum_k D_{\perp f}(k)$, respectively. The nonlinear transfer of entropy $T_f(k)$ is given by Ref. 17

$$\begin{aligned} T_f(k) &= \sum_{k'} \mathcal{T}_f(k, k') \\ &= \sum_{k'} \int d\Theta \frac{T_0}{F_0} f_k^* \left[(k_x - k'_x) \bar{\phi}_{1(k-k')} k'_y f_{k'} \right. \\ &\quad \left. - (k_y - k'_y) \bar{\phi}_{1(k-k')} k'_x f_{k'} \right], \end{aligned}$$

where f_k^* is the complex conjugate of f_k and the sum is over all the k'_x and k'_y . In a similar way, we define a nonlinear transfer term and a spectral density of the electrostatic energy, $T_{\phi}(k)$ and $E_{\phi}(k)$, respectively. Note that the nonlinear term is the only term responsible for a transfer of entropy or of electrostatic energy between different Fourier modes, e.g., between $f_{k'}$ and f_k due to $\bar{\phi}_{1(k-k')}$ (which is linear in $f_{k-k'}$ because of the Poisson equation in Fourier space). This determines a so-called triadic interaction between the modes f_k , $f_{k-k'}$, and $f_{k'}$, which is interpreted as an exchange of energy between modes k and k' due to the property $\mathcal{T}_f(k, k') = -\mathcal{T}_f(k', k)$ (from which follows the conservative behavior of this term, i.e., $\sum_k T(k) = \sum_{k,k'} \mathcal{T}(k, k') = -\sum_{k,k'} \mathcal{T}(k', k) = 0$).

Usually, the nonlinear transfer $T(k)$ is interpreted as (minus) the divergence of a flux in wavenumber space, i.e.,

$$T(k) = - \frac{\partial \Pi(k)}{\partial k},$$

which means that the local energy balance equation has the form

$$\frac{\partial E(k)}{\partial t} + \frac{\partial \Pi(k)}{\partial k} = -D_{\text{tot}}(k), \quad (9)$$

where now D_{tot} represents all the possible dissipation sources. Then, one usually assumes a quasi-stationary state (so $\partial_t E(k) \approx 0$ after a time-average) and considers the inertial range ($D_{\text{tot}}(k) \approx 0$), so the above balance equation reduces to

$$\frac{\partial \Pi(k)}{\partial k} = 0 \quad \Rightarrow \quad \Pi(k) = \Pi_0 = \text{const.},$$

and it can be solved with the help of a closure relation between the flux and the spectrum, if such a relation exists, e.g.,

$$\Pi(k) \sim k^\alpha E(k) \quad \Rightarrow \quad E(k) \propto k^{-\alpha}.$$

However, these arguments apply if the turbulence can be assumed to be in a quasi-stationary state (which is usually the case for driven turbulence) and if there exists a range over which the dissipation and the drive is negligible (i.e., what is commonly called inertial range). Unfortunately, this seems not to be the case for freely decaying sub-Larmor scale turbulence, which is what we are going to study in the present work. Thus, a more general approach based on heuristic and physical arguments is required.

III. SCALINGS AND COLLISIONS

For the almost collisionless (or weakly collisional) case, a scaling theory of the entropy cascade in sub-Larmor scale range has been proposed^{8,9} and it has been tested by means of direct numerical simulations.^{10,11} However, such Kolmogorov-style arguments are based on several assumptions, one of which is the existence of an inertial range in which no dissipation occurs. Nevertheless, gyrokinetics can exhibit multiscale dissipation throughout a wide wave number range, as it was shown in recent papers.^{18–20} With the present work, we want to add a new, simple system to the previously mentioned cases, showing that even in our case multiscale dissipation is present, and moreover, it can also affect the spectra exponents, leading to non-universal power laws. Nonetheless, this work confirms again the weakly collisional theory in the proper limit, extending it in the moderate collisionality limit (i.e., where the spectra still have a significant power law component and they are not just an exponential fall off, which would be the case in the very high collisionality limit).

A. A review of the weakly collisional theory

Before going to the intermediate collisionality case, let us quickly review the weakly collisional theory suggested in Refs. 8 and 9. In our simple system, the only term responsible for the energy transfer among modes is the nonlinear term in Eq. (1), $\mathbf{v}_{\tilde{\phi}} \cdot \nabla F_1$, from which we can readily estimate the nonlinear decorrelation rate ω_{NL} , i.e.,

$$\omega_{NL} \sim k_{\perp} J_0(k_{\perp} \rho) \phi_k k_{\perp} \sim k_{\perp}^{3/2} \phi_k, \quad (10)$$

where we have used the large argument approximation of the Bessel function, $J_0(\zeta) \approx \zeta^{-1/2} \cos(\zeta - \pi/4)$ for $\zeta \gg 1$, since we are interested in the sub-ion-gyroradius scale range cascade, $k_{\perp} \rho \gg 1$. Then, since our aim is to first estimate the

free energy flux, $\omega_{NL} f_k^2 \sim k_{\perp}^{3/2} \phi_k f_k^2$, the next step is to relate the electrostatic potential components ϕ_k to those of the perturbed part of the distribution function f_k . This can be done through the GK Poisson equation, Eq. (2); in the Fourier representation, it reads

$$\phi_k = \tilde{\beta}(k_{\perp}) \int v_{\perp} J_0\left(\frac{k_{\perp} v_{\perp}}{\Omega}\right) \hat{F}_1 dv_{\parallel} dv_{\perp}, \quad (11)$$

where $\tilde{\beta}(k) = 2\pi/(1 - \hat{\Gamma}_0(k)) = 2\pi/(1 - I_0(k^2)e^{-k^2})$, I_0 being the modified Bessel function. In the $k_{\perp} \rho \gg 1$ limit, $\tilde{\beta}(k) \approx \text{const.}$ and the large argument expansion of the Bessel function J_0 gives

$$\phi_k \sim k_{\perp}^{-1/2} \int v_{\perp}^{-1/2} \cos\left(\frac{k_{\perp} v_{\perp}}{\Omega} - \frac{\pi}{4}\right) \hat{F}_1 dv_{\parallel} dv_{\perp}. \quad (12)$$

Now the nonlinear phase mixing argument comes into play and helps us to estimate the above integral by assuming the correspondence of length scales in real space, l , and in perpendicular velocity space,²¹ l_v , due to the nonlinear phase mixing process, i.e., $l_v \sim l$. In fact, let us assume that l is the correlation length of the $\mathbf{E} \times \mathbf{B}$ flow (i.e., of the electrostatic potential ϕ_1). Since in gyrokinetics the particle drift is determined by the fluctuating fields averaged over their gyro-orbits, particles sharing the same gyrocenter position, but having different perpendicular velocities v_{\perp} , are gyro-averaging over different Larmor orbits and thus they are experiencing different drifts in real space. If the difference in the Larmor radii, and thus in $v_{\perp} \propto \rho$, of such orbits is of the order of the correlation length l of the averaged potential, then the two particles are decorrelated and they will perform independent random walks. This, in turns, mean that the distribution function F_1 develops random structures in v_{\perp} -space on the scales $l_v \sim l$ (see Refs. 8 and 9 for further details). Then, because of the random nature, this process, the velocity space integral (12) can be argued to accumulate like a random walk in which the step size scales as l_v and the number of steps scale as l_v^{-1} , so the typical displacement scales as $l_v^{1/2}$. In terms of v_{\perp} -scale conjugate variable p (p is for the Hankel transform in v_{\perp} -space what k_{\perp} is for the Fourier transform in real space), this means

$$\phi_k \sim k_{\perp}^{-1/2} p^{-1/2} f_k \sim k_{\perp}^{-1} f_k, \quad (13)$$

where the last step comes from the nonlinear phase mixing argument $l \sim l_v$ and f_k has to be interpreted⁹ as the root-mean-square value of the fluctuations in \hat{F}_1 , i.e., $f_k \sim \sqrt{\int \hat{F}_1^2 v_{\perp} dv_{\perp} dv_{\parallel}}$. Now, from Eqs. (10) and (13), assuming locality of interactions and constancy of the flux of free energy Π_f , we obtain

$$\omega_{NL} f_k^2 \sim k_{\perp}^{3/2} \phi_k f_k^2 \sim k_{\perp}^{-1/2} f_k^3 \sim \varepsilon_0 = \text{const.}, \quad (14)$$

from which we readily get $f_k \sim \varepsilon_0^{1/3} k_{\perp}^{-1/6}$ and $\phi_k \sim \varepsilon_0^{1/3} k_{\perp}^{-7/6}$ and finally the spectra

$$E_f(k_{\perp}) \sim \varepsilon_0^{2/3} k_{\perp}^{-4/3}, \quad (15a)$$

$$E_\phi(k_\perp) \sim \varepsilon_0^{2/3} k_\perp^{-10/3}. \quad (15b)$$

Note that in the above assumptions, two features can be taken as the main ones, i.e., the nonlinear phase mixing process (i.e., $l \sim l_v$) and the existence of a Kolmogorov-like inertial range over which absolutely no dissipation occurs (i.e., the constancy of the free energy flux $\Pi_f \sim \varepsilon_0 = \text{const}$). In the following, we are going to relax the latter one and to slightly modify the argument from which we are estimating the velocity integral in Eq. (12), while still assuming the nonlinear phase mixing argument $l \sim l_v$ to hold.

B. Taking into account multiscale dissipation

We now extend the above theory in order to take into account collisional effects. First of all, let us assume that the nonlinear phase mixing argument, $l_v \sim l$, is still valid and that the decorrelation rate is still given by the nonlinear term, $\omega_{NL} \sim k_\perp^{3/2} \phi_k$, which is responsible for the conservative energy transfer in wavenumber space. Then, with respect to the weakly collisional case, we argue that the scaling between ϕ_k and f_k becomes steeper due to the collisional smearing of the very small velocity-space scales, i.e.

$$\phi_k \sim k_\perp^{-1-\theta_\nu} f_k, \quad (16)$$

which is estimated from Eq. (12) with the same argument by which the velocity integral accumulates like a random walk, but now the step is a bit larger than in the weakly collisional case, an effect which is represented here via the θ_ν correction. Again, we now have to consider the flux of free energy, i.e., $\omega_{NL} f_k^2 \sim k_\perp^{3/2} \phi_k f_k^2$, together with the scaling in Eq. (16). In this regard, we do not assume a constancy of such flux as in the weakly collisional case (i.e., the existence of an inertial range over which no dissipation occurs), but we allow for a k_\perp -dependent $\Pi_f(k_\perp)$ due to multiscale dissipation

$$\omega_{NL} f_k^2 \sim k_\perp^{3/2} \phi_k f_k^2 \sim k_\perp^{-1/2} f_k^3 \sim \Pi_f(k_\perp). \quad (17)$$

In general, to take into account the dissipation cut-off, the flux can be expressed as a combination of a power law and an exponential, e.g.,

$$\Pi_f(k_\perp) \sim \varepsilon_0 k_\perp^{-\delta_\nu} e^{-\beta k_\perp^\gamma}.$$

Note that, in principle also the exponential parameters depend on the collisionality, i.e., $\beta = \beta_\nu$ and $\gamma = \gamma_\nu$. However, we assume that there exists a k_\perp -range over which the flux is nearly a power law, i.e.,

$$\Pi_f(k_\perp) \approx \varepsilon_0 k_\perp^{-\delta_\nu}. \quad (18)$$

This leads to the following spectra:

$$E_f(k_\perp) \sim \varepsilon_0^{2/3} k_\perp^{-4/3-2(\delta_\nu-\theta_\nu)/3}, \quad (19a)$$

$$E_\phi(k_\perp) \sim \varepsilon_0^{2/3} k_\perp^{-10/3-2(\delta_\nu+2\theta_\nu)/3}, \quad (19b)$$

which are in general steeper than the weakly collisional spectra in Eq. (15), which are however consistently recovered by

the limit $\theta_\nu, \delta_\nu \rightarrow 0$. We finally note that, if we retain the exponential cut-off in the free energy flux $\Pi_f(k_\perp)$, we obtain the same result given in Eq. (19), just multiplied by the exponential cut-off $\exp(-\frac{2}{3}\beta_\nu k_\perp^\gamma)$.

Hereafter, we will refer to the spectra exponents as α_f and α_ϕ , such that $E_f(k_\perp) \propto k_\perp^{-\alpha_f}$ and $E_\phi(k_\perp) \propto k_\perp^{-\alpha_\phi}$.

IV. DIRECT NUMERICAL SIMULATIONS

In order to test these ideas, the nonlinear GK equations, Eqs. (1) and (2), are solved by means of direct numerical simulations with the GENE code in a 4D phase space (x, y, v_\parallel , and μ). The size of the domain in real space is $L_x = L_y = 2\pi\rho$, while the velocity-space domain is bounded by $-3v_T \leq v_\parallel \leq +3v_T$ and $0 \leq \mu \leq 9T_0/B_0$, where $v_T = \sqrt{2T_0/m}$ is the thermal velocity. We use (256, 128, 32, and 96) points in our 4D phase-space and a linearized Landau collision operator acting on F_1/F_0 in gyrocenter coordinates^{22–25} is adopted for $\langle \mathcal{L}[F_1] \rangle$. Note that, since we are using a Fourier representation, the resolution we have indicated as (256, 128) in the Fourier modes actually corresponds to (256, 256) (fully dealiased) grid points in real-space coordinates. We remind the reader that there are no gradients in the background quantities, n_0, T_0 , and B_0 . Thus, the system is initialized with an appropriate perturbed distribution function F_1 , and then it may freely evolve. In the simulations $B_0 = 1$ and $T_0 = 1$, so the μ -grid resolution is high enough to account for the largest k_\perp modes and thus for the expected nonlinear phase mixing argument $l \sim l_v$ to hold also in the numerical framework. We have chosen the same initial condition as given in Ref. 10, i.e., a Maxwellian in velocity space and a sum of $k_x = 2$ and $k_y = 2$ cosines with a small-amplitude white noise χ on all Fourier modes

$$F_1(x, y, v_\parallel, \mu; t = 0) = F_0[\cos(2x/\rho) + \cos(2y/\rho) + \epsilon\chi(x, y)]F_0(v_\parallel, \mu),$$

where F_0 is a constant, ϵ is the (small) amplitude of the white noise with respect to the cosine functions, and F_0 is the background Maxwellian. With time, the system evolves into a turbulent state (see Fig. 1).

Moreover, in order to avoid energy pile-up at the very end of the spectrum (high- k_\perp), an 8th-order k_\perp -hyperdiffusion operator $H_\perp[F_1] = -a_\perp(k_\perp/k_0)^8 F_1$ is added on the RHS of Eq. (1). Note that, due to the very high order of the




FIG. 1. Contour plots of the initial condition n_1 (left) and of a later turbulent state for ϕ_1 (right).

hyperdiffusion and to an appropriate choice of k_0 , $H_\perp[F_1]$ will be relevant only for very high k_\perp and thus negligible in the k_\perp -range of interest for the spectra (e.g., $a_\perp = 0.5$ and $k_0 = 96$ in our runs). We remark that, since we are dealing with decaying turbulence, the spectra must be normalized appropriately at each time step (e.g., w.r.t. the amplitude of the decaying quantity) and then time-averaged over a (collisionality dependent) interval $[t_a, t_b]$. In particular, the parameter t_b can be chosen as the maximum simulation time, $t_{\max} = 120$, for every collision frequency ν , provided that we are in a state of almost completely decayed turbulence (so the spectra do not change if we pass from $t_b = t_{\max}$ to $t_b = t_{\max} - \Delta t$, with Δt sufficiently small). The parameter t_a is collisionality dependent, since it must be chosen in a way such that the turbulence is fully developed, which actually depends on ν . For a quantity $A_{ijs} \equiv A(k_{x,i}, k_{y,j}, t_s)$, the spectrum $E_A(n) = E_A(k_{\perp,n})$ is defined by

$$E_A(n) \equiv \widetilde{\sum}_{(i,j) \in n} E_A(i,j) = \widetilde{\sum}_{ij} \sum_s \frac{w_s |A_{ijs}|^2}{\sum_{i,j} |A_{ijs}|^2},$$

where $\widetilde{\sum}_{(i,j) \in n}$ is the “ring average” over the n -th shell, and, since GENE uses adaptive time steps, w_s are the corresponding “time weights”. In Fig. 2, we report the numerical results for the free energy spectrum $E_f(k_\perp)$ and the electrostatic energy spectrum $E_\phi(k_\perp)$ from simulations with collision frequencies of $\nu = 10^{-6}$, 10^{-5} , and 10^{-4} (black, blue, and

red curves, respectively). In order to make more clear how collisional the turbulence is, according to Refs. 10 and 11, these three cases correspond to a Dorland number of $D = 40$, 527, and 5438, respectively, for $\nu = 10^{-4}$, 10^{-5} , and 10^{-6} . In particular, this justifies the use of hyperdiffusion, since in the very high Dorland number regime, the spectral cutoff $k_{\perp,c} \rho \propto D^{3/5}$ would exceed the resolution; thus, we need to impose an artificial cutoff. Moreover, we used the same hyperdiffusion parameters for all the cases in order to make them comparable. For these cases, the initial time for the average was chosen to be $t_a = 40$, 38, and 35 for $\nu = 10^{-6}$, 10^{-5} , and 10^{-4} , respectively. As we can see, the spectra become steeper with increasing collision frequency ν . This is indeed what can be expected qualitatively from Eq. (19): good agreement with the standard theory, Eq. (15), for low collisionality, and a steepening of the slopes with increasing ν .

We would like to stress that, even though the exponential part of the spectrum becomes more important with increasing collisionality, up to $\nu \sim 10^{-4}$ the power law part is still dominant and the spectrum can be described by a pure power law for an order of magnitude to a very good approximation (see Fig. 2). An overview of the exponents, α_f and α_ϕ , as inferred from the simulation results shown in Fig. 2 via a linear fit is presented in Table I. The uncertainties are estimated from the variation of the exponents due to the choice of the k_\perp -range of fitting, the time window of average and the number of bins for the ring average.

Numerical simulations can also be used to check the hypotheses used in the theory (Sec. III). In particular, we are going to test three fundamental features of the theory: (i) the scaling relation between ϕ_k and f_k , (ii) the free energy flux Π_f , and (iii) the locality of the energy cascade. The first assumption, i.e., the scaling relation in Eq. (16), which reduces to the standard scaling (13) in the low collisionality limit ($\theta_\nu \rightarrow 0$), is displayed in Fig. 3 for the three cases $\nu = 10^{-6}$, 10^{-5} , and 10^{-4} of Fig. 2. Note that in Fig. 3 we have introduced a shift in magnitude of the relation for the three cases for the sake of clarity. The simulations thus demonstrate that the scaling relation between ϕ_k and f_k becomes steeper than k_\perp^{-1} with increasing collisionality, although $\theta_\nu \ll 1$ is only a relatively small correction (e.g., $\theta_\nu \simeq 0.15 \pm 0.05$ for $\nu = 10^{-4}$). This was expected, since we are still in a regime in which the collisions do not affect the non-linear phase mixing argument, $l_p \sim l$, but they just avoid to form a lot of small scale structures in v_\perp -space.

In Fig. 4, the second assumption, regarding the flux of free energy, Eq. (18), is investigated. Here, the flux is normalized to the total dissipation. The non-constancy of the flux reflects the fact that dissipation is actually effective at all scales, since for purely conservative spectral energy




FIG. 2. $E_f(k_\perp)$ (top) and $E_\phi(k_\perp)$ (bottom) for three different collision frequencies: $\nu = 10^{-6}$, 10^{-5} , and 10^{-4} (black squares, blue diamonds, and red circles, respectively). The dashed lines represent the weakly collisional theory.

TABLE I. Spectral exponents extracted from the simulations shown in Fig. 2.

ν	α_f	α_ϕ
10^{-6}	1.35 ± 0.05	3.45 ± 0.10
10^{-5}	1.50 ± 0.05	3.75 ± 0.10
10^{-4}	2.05 ± 0.15	4.45 ± 0.25




FIG. 3. Scaling relation ϕ_k/f_k for three different collision frequencies: $\nu = 10^{-6}$, 10^{-5} , and 10^{-4} (black squares, blue triangles, and red diamonds, respectively). The dashed line corresponds to the scaling k_{\perp}^{-1} predicted by Ref. 9.

transfer (i.e., the standard picture of the inertial range), one expects $\Pi_f/D_{\text{tot}} = 1$ for an extended region in k -space, until the dissipation range begins and an exponential fall-off appears. However, this is only approximately the case even at the lowest collisionality, $\nu = 10^{-6}$, for which $(\Pi_f/D_{\text{tot}})_{\text{max}} \sim 0.95$. Meanwhile, we obtain $(\Pi_f/D_{\text{tot}})_{\text{max}} \sim 0.75$ for $\nu = 10^{-5}$ and $(\Pi_f/D_{\text{tot}})_{\text{max}} \sim 0.5$ for $\nu = 10^{-4}$, after which the flux is not even constant (it can be considered a power law, as a first approximation). This behavior clearly deviates from the standard picture of an inertial range.

Moreover, the differences between the assumptions made in the standard theory and the fluxes in Fig. 4 are way more evident than the deviations found in the scalings (Fig. 3). In other words, δ_{ν} is deviating from zero more rapidly due to the multiscale dissipation: e.g., we estimate $\delta_{\nu} \simeq 1.0 \pm 0.2$ for $\nu = 10^{-4}$. Again, this means that the weaker point is to assume the standard picture of an inertial range bridging the injection and the dissipation scales: this is something that was already observed in gyrokinetics for other type of systems (see Refs. 18–20). As a final remark on these two assumptions, we note that not only qualitative agreement is found: the values of θ_{ν} and δ_{ν} estimated from simulations are also in quantitative agreement with the fitted exponents of the spectra. In fact, for instance, we found




FIG. 4. Normalized free energy flux $\Pi_f(k_{\perp})/D_{\text{tot}}$ for $\nu = 10^{-6}$, 10^{-5} , and 10^{-4} (black squares, blue triangles, and red circles, respectively). The green dashed line corresponds to unity, while the other colored dashed lines corresponds to the maximum of the respective flux.




FIG. 5. Shell-to-shell transfer T_n^m for $\nu = 10^{-6}$ (right) and for $\nu = 10^{-4}$ (left).

$\alpha_f \simeq 2.05 \pm 0.15$ and $\alpha_{\phi} \simeq 4.45 \pm 0.25$ for the $\nu = 10^{-4}$ case (Table I), while the values of θ_{ν} and δ_{ν} extracted from the simulations (Figs. 3 and 4) predict $\alpha_f \simeq 1.90 \pm 0.15$ and $\alpha_{\phi} \simeq 4.20 \pm 0.25$. Considering the approximations made and the uncertainties, this is a good agreement.

We finally test the third assumption, i.e., the locality of the energy cascade. This feature can be checked by looking at the shell-to-shell transfer, T_n^m , i.e., the energy exchange between the k_{\perp} shells. Defining the shells as $\{k_n\}_{n=1,2,\dots,N}$, where we fix three parameters, k_a and k_b and A , such that $k_1 = k_a$, $k_2 = k_a + k_b$, and $k_n = (k_a + k_b)2^{(n-2)/A} = k_2 2^{(n-2)/A}$ for $n = 2, \dots, N$. In the following, $k_a = k_b = 4$ and $A = 5$ will be adopted. Note that T_n^m is the discrete version of $\mathcal{T}_f(k', k)$. The shell-to-shell transfer for $\nu = 10^{-6}$ and $\nu = 10^{-4}$ is shown in Fig. 5 (left and right panel, respectively): we can see that the energy transfer is very local, i.e., the free energy exchange is relevant only between neighboring shells, thus verifying the locality assumption. Moreover, the antisymmetry of T_n^m is immediately evident from the plot, which means it is a non-dissipative term that only transfers energy between the modes, as to be expected. From Fig. 5, we also recognize the features already seen in Fig. 4 about the free energy flux $\Pi_f(k_{\perp})$, i.e., the non-constancy of the flux for $\nu = 10^{-4}$, denoted by the softening of the colors, in contrast to its nearly constant behavior for the $\nu = 10^{-6}$ case.

V. CONCLUSIONS AND DISCUSSION

We have presented a theoretical and numerical study of freely decaying electrostatic turbulence in the framework of collisional gyrokinetic theory of magnetized plasmas. A reduced 4D (2D2V) phase space $(x, y, v_{\parallel}, \mu)$ is considered: the 2D real space is perpendicular to the background magnetic field, thus avoiding parallel effects such as the Landau damping to be effective. No background gradients were considered, making the system as simple as possible, and thus focusing on the two effects of interest: the $E \times B$ nonlinearity and collisions. The nonlinear term introduces a perpendicular (nonlinear) phase mixing,^{8,9} causing the perturbed part of the distribution function to develop structures in v_{\perp} -space, which are related to those in real space (or, equivalently, to k_{\perp}). In particular, the higher the k_{\perp} is, the finer those v_{\perp} -structures are. However, the relation between v_{\perp} -scales and k_{\perp} is affected by the collisionality of the system, which in practice has the function of limiting the finest v_{\perp} -scales. Then, due to the relationship between v_{\perp} scales and real-space scales, this

has a direct influence on the k_{\perp} scalings between ϕ_k and f_k . This was indeed shown by means of direct numerical simulations, even if it remains a small effect, provided that the collisionality is not too high. In addition, allowing for the presence of multiscale dissipation in sub-Larmor scale fluctuations has immediate consequences on one of the very fundamental assumptions made by standard Kolmogorov-like theories, i.e., on the existence of an inertial range. In fact, regarding this point, we have shown that dissipation occurs at all scales, regardless of the collisionality regime and, moreover, that for intermediate collisionality the free energy flux is not even nearly constant anymore. This leaves a very important fingerprint on the spectra, making in fact the power law collisionality dependent and thus allowing for non-universal power laws.

Despite the relative simplicity of the system under study here, it seems plausible that these results can be generalized and applied to more complicated systems. For instance, it is reasonable to expect that also 3D GK turbulence in toroidal fusion devices can exhibit variable power law scalings depending on the parameter settings. Such behaviour has indeed been observed before, and the present work offers a possible explanation. Similarly, 3D GK turbulence simulations applied to the solar wind dissipation range display non-universal power law scalings as well as exponential corrections. Again, the present work may provide a key to the understanding of this effect. Follow-up studies will have to clarify if these conjectures are correct.

ACKNOWLEDGMENTS

We would like to thank H. Doerk, V. Bratanov, H. Sceats, G. G. Plunk, and A. A. Schekochihin for very useful discussions. The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC Grant Agreement No. 277870. This project

has received funding from the Euratom research and training programme 2014–2018.

- ¹U. Frisch, *Turbulence* (Cambridge University Press, 1995).
- ²G. Falkovich and K. R. Sreenivasan, *Phys. Today* **59**(4), 43 (2006).
- ³E. A. Friemann and L. Chen, *Phys. Fluids* **25**, 502 (1982).
- ⁴A. J. Brizard and T. S. Hahm, *Rev. Mod. Phys.* **79**, 421 (2007).
- ⁵J. A. Krommes, *Annu. Rev. Fluid Mech.* **44**, 175 (2012).
- ⁶A. A. Schekochihin, S. C. Cowley, W. Dorland, G. W. Hammett, G. G. Howes, E. Quataert, and T. Tatsuno, *ApJS* **182**, 310 (2009).
- ⁷A. Bañón Navarro, P. Morel, M. Albrecht-Marc, D. Carati, F. Merz, T. Görler, and F. Jenko, *Phys. Rev. Lett.* **106**, 055001 (2011).
- ⁸A. A. Schekochihin, S. C. Cowley, W. Dorland, G. W. Hammett, G. G. Howes, G. G. Plunk, E. Quataert, and T. Tatsuno, *Plasma Phys. Controlled Fusion* **50**, 124024 (2008).
- ⁹G. G. Plunk, S. C. Cowley, A. A. Schekochihin, and T. Tatsuno, *J. Fluid Mech.* **664**, 407 (2010).
- ¹⁰T. Tatsuno, W. Dorland, A. A. Schekochihin, G. G. Plunk, M. Barnes, S. C. Cowley, and G. G. Howes, *Phys. Rev. Lett.* **103**, 015003 (2009).
- ¹¹T. Tatsuno, M. Barnes, S. C. Cowley, G. G. Howes, R. Numata, G. G. Plunk, and A. A. Schekochihin, *J. Plasma Fusion Res. SERIES* **9**, 509 (2010).
- ¹²F. Jenko, W. Dorland, M. Kotschenreuther, and B. N. Rogers, *Phys. Plasmas* **7**, 1904 (2000).
- ¹³T. Dannert and F. Jenko, *Phys. Plasmas* **12**, 072309 (2005).
- ¹⁴T. Görler, X. Lapillonne, S. Brunner, T. Dannert, F. Jenko, F. Merz, and D. Told, *J. Comput. Phys.* **230**, 7053 (2011).
- ¹⁵L. D. Landau, *J. Phys. (USSR)* **10**, 25 (1946).
- ¹⁶G. W. Hammett, W. Dorland, and F. W. Perkins, *Phys. Fluids B* **4**, 2052 (1992).
- ¹⁷A. Bañón Navarro, P. Morel, M. Albrecht-Marc, D. Carati, F. Merz, T. Görler, and F. Jenko, *Phys. Plasmas* **18**, 092303 (2011).
- ¹⁸B. Teaca, A. Bañón Navarro, F. Jenko, S. Brunner, and L. Villard, *Phys. Rev. Lett.* **109**, 235003 (2012).
- ¹⁹D. R. Hatch, F. Jenko, A. Bañón Navarro, and V. Bratanov, *Phys. Rev. Lett.* **111**, 175001 (2013).
- ²⁰B. Teaca, A. Bañón Navarro, and F. Jenko, “The energetic coupling of scales in gyrokinetic plasma turbulence,” *Phys. Plasmas* **21**, 072308 (2014).
- ²¹If δv_{\perp} is the typical velocity scale length between the fluctuations developed by the distribution function F_1 in v_{\perp} -space (see later in the text), then we can define $l_r \sim (\delta v_{\perp}/v_T)\rho$ (see, e.g., Ref. 8).
- ²²F. Merz, Ph.D. dissertation (Universität Münster, 2008).
- ²³I. G. Abel, M. Barnes, S. C. Cowley, W. Dorland, and A. A. Schekochihin, *Phys. Plasmas* **15**, 122509 (2008).
- ²⁴M. Barnes, I. G. Abel, T. Tatsuno, A. A. Schekochihin, S. C. Cowley, and W. Dorland, *Phys. Plasmas* **16**, 072107 (2009).
- ²⁵H. Doerk, Ph.D. dissertation (Universität Ulm, 2012).